



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2014

Woody plant phylogenetic diversity mediates bottom-up control of arthropod biomass in species-rich forests

Schuldt, Andreas ; Baruffol, Martin ; Bruelheide, Helge ; Chen, Simon ; Chi, Xiulian ; Wall, Marcus ;
Assmann, Thorsten

Abstract: Global change is predicted to cause non-random species loss in plant communities, with consequences for ecosystem functioning. However, beyond the simple effects of plant species richness, little is known about how plant diversity and its loss influence higher trophic levels, which are crucial to the functioning of many species-rich ecosystems. We analyzed to what extent woody plant phylogenetic diversity and species richness contribute to explaining the biomass and abundance of herbivorous and predatory arthropods in a species-rich forest in subtropical China. The biomass and abundance of leaf-chewing herbivores, and the biomass dispersion of herbivores within plots, increased with woody plant phylogenetic diversity. Woody plant species richness had much weaker effects on arthropods, but interacted with plant phylogenetic diversity to negatively affect the ratio of predator to herbivore biomass. Overall, our results point to a strong bottom-up control of functionally important herbivores mediated particularly by plant phylogenetic diversity, but do not support the general expectation that top-down predator effects increase with plant diversity. The observed effects appear to be driven primarily by increasing resource diversity rather than diversity-dependent primary productivity, as the latter did not affect arthropods. The strong effects of plant phylogenetic diversity and the overall weaker effects of plant species richness show that the diversity-dependence of ecosystem processes and interactions across trophic levels can depend fundamentally on non-random species associations. This has important implications for the regulation of ecosystem functions via trophic interaction pathways and for the way species loss may impact these pathways in species-rich forests.

DOI: <https://doi.org/10.1007/s00442-014-3006-7>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-98301>

Journal Article

Originally published at:

Schuldt, Andreas; Baruffol, Martin; Bruelheide, Helge; Chen, Simon; Chi, Xiulian; Wall, Marcus; Assmann, Thorsten (2014). Woody plant phylogenetic diversity mediates bottom-up control of arthropod biomass in species-rich forests. *Oecologia*, 176(1):171-182.

DOI: <https://doi.org/10.1007/s00442-014-3006-7>

Woody plant phylogenetic diversity mediates bottom-up control of arthropod biomass in species-rich forests

Andreas Schuldt^{1,*}, Martin Baruffol², Helge Bruelheide^{3,4}, Simon Chen⁵, Xiulian Chi⁶, Marcus Wall⁷, Thorsten Assmann¹

¹Leuphana University Lüneburg, Institute of Ecology, Scharnhorststr. 1, D-21335 Lüneburg, Germany; ²University of Zurich, Institute of Evolutionary Biology and Environmental Studies, Winterthurerstr. 190, CH-8057 Zurich, Switzerland; ³University of Halle, Institute of Biology/Geobotany and Botanical Garden, Am Kirchtor 1, D-06108 Halle, Germany; ⁴German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Deutscher Platz 5e, D-04103 Leipzig, Germany; ⁵Magdalene College, GB-CB3 0AG Cambridge; ⁶Department of Ecology, College of Urban and Environmental Sciences, and Key Laboratory for Earth Surface Processes of Ministry of Education, Peking University, Beijing, China; ⁷University of Halle, Natural History Collections, Domplatz 4, D-06108 Halle

* Correspondence: schuldt@uni.leuphana.de; Tel.: +49 4131 6772965; Fax: +49 4131 6772808

Running head: Plant phylodiversity affects arthropods

Abstract

Global change is predicted to cause non-random species loss in plant communities, with consequences for ecosystem functioning. However, beyond the simple effects of plant species richness, little is known about how plant diversity and its loss influence higher trophic levels, which are crucial to the functioning of many species-rich ecosystems. We analyzed to what extent woody plant phylogenetic diversity and species richness contribute to explaining the biomass and abundance of herbivorous and predatory arthropods in a species-rich forest in subtropical China. The biomass and abundance of leaf-chewing herbivores, and the biomass dispersion of herbivores within plots, increased with woody plant phylogenetic diversity. Woody plant species richness had much weaker effects on arthropods, but interacted with plant phylogenetic diversity to negatively affect the ratio of predator to herbivore biomass. Overall, our results point to a strong bottom-up control of functionally important herbivores mediated particularly by plant phylogenetic diversity, but do not support the general expectation that top-down predator effects increase with plant diversity. The observed effects appear to be driven primarily by increasing resource diversity rather than diversity-dependent primary productivity, as the latter did not affect arthropods. The strong effects of plant phylogenetic diversity and the overall weaker effects of plant species richness show that the diversity-dependence of ecosystem processes and interactions across trophic levels can depend fundamentally on non-random species associations. This has important implications for the regulation of ecosystem functions via trophic interaction pathways and for the way species loss may impact these pathways in species-rich forests.

Keywords: BEF China; biodiversity; ecosystem function; herbivores; predators; species richness; trophic interactions

Introduction

The diversity of life on earth strongly influences the dynamics and properties of ecosystems, and global change-induced biodiversity loss may thus significantly alter ecosystem functioning and service provisioning (Hooper et al. 2012; Naeem et al. 2012). Most notably, plant species loss has been shown to negatively affect a wide range of ecosystem functions, such as biomass production and carbon sequestration (Cardinale et al. 2011; Cardinale et al. 2012). However, recent work has also indicated that plant species richness impacts on the abundance and diversity of higher trophic levels (e.g. Unsicker et al. 2006; Haddad et al. 2009; Sobek et al. 2009b; Scherber et al. 2010). This has important ecological consequences, as diversity-dependent effects on herbivores and predators can feed back on the producer level and strongly mediate biodiversity-ecosystem function relationships under real-world conditions (Haddad et al. 2009; Schuldt et al. 2010).

However, we are far from having a general understanding of such diversity-dependent trophic interaction effects (see also Cardinale et al. 2011), as even basic relationships between plant diversity and the diversity or abundance of herbivores and predators often seem not to show a consistent pattern (Koricheva et al. 2000; Vehviläinen et al. 2007; Sobek et al. 2009a; Castagneyrol and Jactel 2012; Schuldt et al. 2014b). While such inconsistent patterns potentially indicate relevant biological differences among systems, their interpretation is complicated by two important aspects. First of all, many studies are biased toward analyzing relatively species-poor and often simplified plant communities, in which it may be difficult to distinguish the effects of community composition from diversity effects (Nadrowski et al. 2010). Extrapolating to more diverse systems is hardly possible. Yet, information on species-rich systems may be particularly crucial, as diversity-dependent interactions between plants and higher trophic levels can play a key role in the functioning of such systems (e.g. Terborgh 2012). Secondly, most studies have focused on the effects of plant species richness as a very basic measure of biodiversity. However, relationships with herbivores or predators might be

more complex and not necessarily captured well by plant species richness alone. Plant species loss in natural communities often occurs, and may cascade to affect higher trophic levels, in a non-random way (Srivastava and Vellend 2005; Thebault et al. 2007; Cavender-Bares et al. 2009). Species may get lost in a phylogenetically structured manner that reflects phylogenetically conserved functional adaptations to their biotic and abiotic environments, such as the phylogenetically structured host selection of many consumers (e.g. Weiblen et al. 2006). This could explain why, even in plant species-rich systems, plant community composition was repeatedly found to be a better predictor of herbivore and predator assemblage structure than species richness (Perner et al. 2005; Rzanny et al. 2013). Measures of diversity that incorporate the relatedness among species in terms of evolutionary and functional similarity may thus substantially improve our understanding of diversity relationships across trophic levels (Dinnage et al. 2012; Pellissier et al. 2013).

Plant phylogenetic diversity qualifies as a particularly comprehensive predictor in this respect, as it may not only account for complex functional trait interactions that affect higher trophic levels (if key functional traits are phylogenetically conserved; Cavender-Bares et al. 2009; Srivastava et al. 2012), but can also indicate the effects of shared evolutionary and biogeographic history among species (Futuyma and Agrawal 2009; Pellissier et al. 2013). Dinnage et al. (2012) recently showed in a grassland experiment that plant phylogenetic diversity strongly interacted with plant species richness to affect arthropod diversity, with effects of plant species richness becoming stronger at high levels of plant phylogenetic diversity. For natural systems, however, where processes of non-random community assembly may strongly affect the phylogenetic structure of plant and animal communities, the effects of phylogenetic diversity on arthropods remain poorly explored. This applies particularly to species-rich subtropical and tropical forests, where arthropods play an important role in maintaining the high tree diversity and may contribute to structuring plant phylogenetic community composition (Cavender-Bares et al. 2009; Paine et al. 2012; Terborgh 2012).

Here, we analyze to what extent the phylogenetic diversity and species richness of woody plants contribute to explaining the biomass and abundance of herbivorous and predatory arthropods along gradients of woody plant species richness and stand age in a highly diverse subtropical forest in China. Biomass, in particular, is directly related to the functional impact of consumers (Saint-Germain et al. 2007; Reiss et al. 2011), but its relationship with plant diversity in such highly diverse forests is poorly understood (and the same even applies to the more frequently studied abundance of arthropods; Whitfeld et al. 2012). We focus on the overall biomass and abundance of predators, leaf chewing herbivores and sucking herbivores, which allows us to obtain insight into the net ecosystem effect of key functional groups in this system. In addition, we test for diversity effects on the variability in biomass distributions among individuals in each of our 27 study plots, which might be related to the degree of resource differentiation in arthropod assemblages (Rudolf 2012).

Plant diversity might have a direct effect on herbivore and predator assemblages via increased and more stable resource diversity, with positive effects on generalist herbivores and predators (the ‘dietary-mixing’ and ‘enemies’ hypotheses; Root 1973; Haddad et al. 2009; Dinnage 2013) and negative effects on specialized herbivores (the ‘resource-concentration’ hypothesis; Root 1973; but see Plath et al. 2012). It might also promote consumers indirectly via diversity-dependent effects of plant productivity on consumer biomass (the ‘more-individuals’ hypothesis; Srivastava and Lawton 1998). We hypothesize that (i) the biomass (and its variability within study plots) and abundances of both herbivores and predators increase with woody plant diversity in our study system. Specifically, we expect that (ii) changes in biomass and abundance are better predicted by plant phylogenetic diversity than plant species richness, as phylogenetic diversity better explains the complexity of evolutionary and functional characteristics that may underlie diversity effects. As woody plant diversity in our study plots has been found to increase primary productivity (Baruffol et al. 2013), we include alternative analyses that substitute productivity data for diversity metrics to

test whether (iii) resource diversity or plant primary productivity underlie potential diversity effects. As support for both the resource diversity and productivity mechanism has been controversial, even for the much better studied grassland systems (Haddad et al. 2009; Borer et al. 2012), knowledge of these potential impacts of these mechanisms on herbivores and predators in highly diverse forests will advance our general understanding of the community-level consequences of changes in biodiversity.

Materials and methods

Study site and plot design

The study was conducted in the Gutianshan National Nature Reserve (29°14' N; 118°07' E) in Zhejiang province, south-east China. The reserve covers 8000 ha of evergreen mixed broadleaved forest on a sloping terrain (300-1260 m a.s.l.). In the reserve, about 1430 seed plant species, 260 of them woody, have been recorded (Legendre et al. 2009; Bruehlheide et al. 2011). The subtropical monsoon climate is characterized by a mean annual temperature of 15.3°C and a mean annual precipitation of about 2000 mm (Hu and Yu 2008), with most of the rainfall occurring in May and June (Geißler et al. 2012).

As part of the BEF (Biodiversity and Ecosystem Functioning) China project (www.bef-china.de), 27 study plots of 30 x 30 m were established in 2008, which were randomly spread (as far as logistically feasible; distance among plots was on average 2.6 km \pm 2.3 SD (range 0.1 - 8.6 km)) across the reserve. Plot selection followed a stratified sampling design based on woody plant species richness (25-69 species per plot) and stand age (with < 20 to > 80 years since the last logging events, or, in some cases, since agricultural activities). Details of plot selection and plot characteristics (including data on the plant species composition of all 27 plots) are given in Bruehlheide et al. (2011).

Arthropod sampling

Arthropods were sampled at three time periods to capture seasonal patterns in the herbivore assemblages, i.e. September/October 2011 (toward the end of the growing season), April 2012 (before the rainy season, after the start of the growing season) and June 2012 (at the end of the rainy season, peak of the growing season). We used a beating technique that allows for a direct assessment of arthropods at the level of individual trees (Ødegaard et al. 2005; Campos et al. 2006; Wardhaugh et al. 2012). Arthropods were knocked down from 25 tree and shrub saplings onto a beating sheet (70 cm diameter). Saplings were selected every two meters along a transect running diagonally through the plot. The composition of these 25 randomly selected saplings very well mirrored the differences in the community composition among plots of the overall woody plant communities (Procrustes correlation = 0.90; $P < 0.001$; see Methods in Electronic Supplementary Material (ESM)). Sapling species identity was determined with the help of local experts. The height of each sampled sapling was recorded in the field. Mean height was 1.77 m, with a SD of 0.48 m, indicating that a similar volume of plant structures was sampled from above the beating sheet for most individuals (see also Campos et al. 2006). However, to account for potential effects of tree height on arthropod samples, individual sapling height was included as a covariate in the statistical analyses. Transect lines varied between the fall and spring surveys, thus sampling a different set of tree and shrub individuals, but were identical between the spring and summer surveys for logistical reasons.

Arthropods were sorted to higher taxa (basically order level, but with further distinctions in orders such as Coleoptera, Hymenoptera, Hemiptera, to distinguish predators and different herbivore guilds; see Table S1 in ESM for details) and classified as (mainly) predators or herbivores (following the classification of Novotny et al. 2010 and using morphological characteristics such as mandible shapes where necessary). For the herbivores, we distinguished between leaf chewers and sap suckers; other herbivore guilds were too rare for separate analysis. As our main focus was on overall biomass and abundance patterns, we

176 did not sort arthropods to (morpho)species, which can be challenging and error-prone for
177 highly diverse taxa (e.g. many juvenile spiders and lepidopteran caterpillars; e.g.
178 Strutzenberger et al. 2011). Biomass and abundance patterns are particularly important in
179 determining the functional impact of primary and secondary consumers (Saint-Germain et al.
180 2007; Reiss et al. 2011), and many studies have shown that overall patterns at the level of
181 important functional groups can be highly informative for our understanding of the regulation
182 of ecosystem functions in light of changing plant diversity (Perner et al. 2005; Vehviläinen et
183 al. 2007; Vehviläinen et al. 2008; Haddad et al. 2009; Borer et al. 2012; Whitfield et al. 2012).
184 For all predatory and herbivorous arthropods, body length (excluding body appendages such
185 as ovipositors and antennae) was measured to the closest 0.1 mm under a stereomicroscope
186 with a built-in micrometer gauge. Biomass for each sampled individual was estimated on the
187 basis of taxon-specific body length-biomass equations of Hóðar (1996) and Wardhaugh
188 (2013).

190 *Environmental data*

191 Woody plant species richness for each plot was based on a complete inventory of all tree and
192 shrub individuals > 1 m height, measured at the time of plot establishment in 2008. Data for
193 the calculation of the phylogenetic diversity of the woody plant communities (see statistics)
194 was obtained from an ultrametric phylogenetic tree of all angiosperm woody species recorded
195 in the 27 study plots (Purschke et al. 2014). The tree was based on rbcL and matK sequences
196 of the species (downloaded from NCBI Genbank; <http://www.ncbi.nlm.nih.gov>), which were
197 aligned in Bioedit and processed in MEGA5 (Tamura et al. 2011) to obtain a phylogenetic
198 tree based on maximum likelihood (ML). The ultrametric tree was computed from the ML
199 topology using penalized likelihood, with branch lengths indicating divergence time.

200 Plot age was estimated from tree stem cores and diameter at breast height
201 measurements (to the closest 0.1 mm) (Bruehlheide et al. 2011). Plot age was correlated with,

and used as a comprehensive measure of, plot characteristics that change with succession, such as canopy cover (Pearson's $r = 0.72$; $P < 0.001$ for the correlation with plot age) and total basal area of woody plants ($r = 0.76$; $P < 0.001$) (see Schuldt et al. 2010). We also accounted for the topographic variability of our study site, which may further affect environmental plot conditions, by including the elevation (m) of the plots in our analyses.

Plot-level primary productivity over a two year period was inferred from tree growth data in 2008 and 2010. Basal area increments of all trees > 10 cm diameter at breast height, assessed with permanent dendrometer bands or with measuring tapes, were calculated and used as a proxy for relative tree growth (see Baruffol et al. 2013 for details). As a number of trees were destroyed in two plots due to illegal harvesting before measurements in 2010, productivity data was only available for 25 plots.

Statistical analysis

Woody plant phylogenetic diversity (Q_{phyl}) was calculated as the abundance-weighted version of Rao's quadratic entropy Q (Botta-Dukát 2005). Q_{phyl} was not completely independent of woody plant species richness ($r = 0.48$; $P = 0.011$) and plot age ($r = 0.57$; $P = 0.002$). As a metric of phylogenetic diversity that reflects the extent to which woody plant communities are phylogenetically more clustered or overdispersed than expected by chance, we thus calculated standardized effect sizes of Q_{phyl} (Q_{phyl} s.e.s.) based on the null model '1s' in Hardy et al. (2008). These s.e.s. values are independent of a given plant community's species richness ($r = -0.26$; $P = 0.186$ in our study). In our case, they were also unrelated to plot age ($r = -0.01$; $P = 0.959$). Standardized effect sizes were calculated as the observed phylogenetic diversity relative to expected values from the random communities: $\text{ses} = (\text{observed phylogenetic diversity index score} - \text{mean expected index score}) / \text{standard deviation of the index}$ (Gotelli and Rohde 2002).

Based on Rao's Q , we also quantified the functional dispersion of the biomass of all

leaf chewer, sap sucker or predator individuals per plot and sampling time (Q_{Bio}). Higher values of Q_{Bio} indicate larger biomass dissimilarity of the individuals of a given group within a study plot, which may point to higher resource differentiation among individuals (Schleuter et al. 2010). Q_{Bio} was not dependent on the number of individuals in a given plot (chewers: $r = -0.07$; $P = 0.564$; suckers: $r = -0.01$; $P = 0.913$; predators: $r = -0.19$; $P = 0.084$). Calculations of Rao's Q were based on standardized variables (mean = 0, SD = 1) and a Euclidean species distance matrix. All continuous predictors (i.e. all variables except sampling time) were standardized prior to the analysis. Correlations among predictors (all with an $r \leq 0.57$) indicated that none of the predictors were highly collinear. The mean biomass and abundances of leaf-chewing herbivores, sap-sucking herbivores, predatory arthropods, and the predator-herbivore biomass and abundance ratios per plot and sampling time—as well as the functional dispersion (Q_{Bio}) of leaf chewer, sucker and predator biomass—were used as response variables. Arthropod biomass and abundance values were averaged for each of the three sampling times across the 25 saplings sampled per plot as different tree and shrub individuals were included in the three different sampling campaigns (see above). Linear mixed effects models were used to account for potential effects of temporal or spatial pseudoreplication. Plot identity was included as a random effect. We also tested for a random interaction effect of sampling time and plot identity, but likelihood ratio tests indicated that this term was not significant and could be dropped. As fixed effects, we included sampling time, elevation, sapling height, plot age, woody plant species richness, woody plant phylogenetic diversity (standardized effect sizes Q_{phyl} s.e.s), as well as all two-way interactions between sampling time, plot age, species richness and phylogenetic diversity. The number of woody plant species sampled in a plot had no effect on any of the response variables and was not included in the models. The response variables and woody plant species richness were log-transformed to improve model fit. The full models with all predictors (i.e. in the form of: response ~ sampling time + elevation + sapling height + plot age + woody plant species richness +

woody plant phylogenetic diversity (Q_{phyl} s.e.s.) + time:age + time:richness +
time:phylodiversity + age:richness + age:phylodiversity + richness:phylodiversity ,
random= ~ 1 |plot) were simplified by excluding predictor variables in an automated stepwise
procedure based on the AICc (Burnham and Anderson 2004) and maximum likelihood
estimation. The models with the smallest number of predictors and the lowest global AICc
were chosen as the most parsimonious, best-fit models for each response variable. Model
residuals were checked for normality and homogeneity of variances. To assess whether
potential effects of tree diversity could be explained by plot-level primary productivity, we re-
ran all analyses that indicated diversity effects in the minimal models, replacing the measures
of diversity (woody plant species richness and phylogenetic diversity) by plot-level
productivity data. Productivity in the study plots was previously found to be strongly related
to woody plant species and phylogenetic diversity (Baruffol et al. 2013) and, to avoid
statistical biases due to collinearity and to keep model complexity to an acceptable level, we
did not directly include productivity in the models which tested for diversity effects. All
analyses were conducted in R 3.1.0 (<http://www.R-project.org>) with the packages *picante*
(Pinheiro et al. 2014) and *nlme* (Kembel et al. 2010).

Results

In total, we recorded 6950 arthropods with a total biomass of 29,167 mg across the three
sampling periods. Predators were most abundant (4737 individuals; 79% spiders), followed by
leaf-chewing herbivores (1282 individuals; 41% lepidopteran caterpillars, 32% orthopterans)
and sap-suckers (931 individuals; 74% Auchenorrhyncha). However, biomass was higher for
leaf chewers (16,730 mg; 42% orthopterans, 37% lepidopteran caterpillars) than for predators
(9387 mg; 66% spiders) and sap suckers (3050 mg; 82% Auchenorrhyncha). Biomass and
abundance correlated particularly strongly for sap suckers (Pearson's $r = 0.74$; $P < 0.001$) but
less so for predators ($r = 0.38$; $P < 0.001$) and leaf chewers ($r = 0.37$; $P < 0.001$). Predator and

herbivore biomass were not significantly correlated ($P > 0.80$ in all cases). In contrast, predator abundance was positively related to leaf chewer abundance ($\beta = 0.26 \pm 0.09 \text{ SE}$; $t = 2.85$; $P = 0.006$ for log-transformed abundance values in a mixed model including sampling time as a covariate) and tended to slightly increase with the abundance of sap suckers ($\beta = 0.07 \pm 0.05 \text{ SE}$; $t = 1.37$; $P = 0.175$).

In general, mixed models for both biomass and abundance pointed to similar variables affecting patterns within the predator and the two herbivore groups across the 27 study plots. However, diversity effects were always more pronounced for biomass as compared to mere abundance patterns (Table 1, Table S2). Thus, in the following we focus on biomass patterns (see Table S2 in ESM for abundance patterns). Leaf chewer biomass was particularly strongly affected by, and increased with, woody plant phylogenetic diversity (Q_{phyl} s.e.s.) (Table 1, Fig. 1a). Leaf chewer biomass further tended to increase with woody plant species richness, but this effect was not significant ($P = 0.07$; Table 1). In contrast to leaf chewers, sap-sucking herbivores were not affected by plant phylogenetic diversity, showing only a response to sampling period (Table 1). Likewise, the biomass of predators was not related to plant phylogenetic diversity, but increased with plot age (Fig. 1b). In contrast, the ratio of predator to herbivore biomass was strongly affected by the interaction between woody plant phylogenetic diversity and plant species richness (a pattern that was not detectable with mere abundance data; Table S2). The ratio of predators to herbivores was highest in plant species-poor plots with lower than expected phylogenetic diversity and strongly decreased with increasing plant phylogenetic diversity and plant species richness (Fig. 1c).

The dissimilarity in biomass among leaf chewer individuals within plots—measured as biomass dispersion Q_{Bio} —was lowest in plant species-poor plots with lower than expected phylogenetic diversity and strongly increased with both woody plant phylogenetic diversity (Q_{phyl} s.e.s.) and plant species richness (Table 2, Fig. 2). Sap sucker biomass dispersion

showed a response to plant phylogenetic diversity only during summer (Time 3; Table 2). Biomass dispersion among predators was not related to plant phlogenetic diversity, but decreased with woody plant species richness (only significant at Time 2; Table 2).

The effects of plant phylogenetic diversity on leaf chewing herbivores and the predator-herbivore ratio were also evident when observed plant phylogenetic diversity instead of standardized effect sizes were analyzed (however, with changes in the roles of plot age and plant species richness due to the correlation of these variables with observed phylogenetic diversity; Tables S5 and S6, Fig. S1).

Plot-level primary productivity did not underlie the effects of plant phylogenetic diversity on arthropods. Plant productivity was not retained in any of the minimal models when the analyses showing significant effects of Q_{phyl} s.e.s. were re-run with measures of woody plant diversity replaced by plant productivity (Tables S3 and S4).

Discussion

Our study highlights how effects of plant diversity that go beyond simple effects of increasing species numbers contribute to controlling consumer biomass across trophic levels in a highly diverse forest. Leaf chewer biomass and the biomass dispersion of herbivores strongly increased across forest stands with increasing plant phylogenetic diversity. In contrast, effects of woody plant species richness were less frequent and less pronounced, interacting with plant phylogenetic diversity to affect arthropods in some cases. Overall, our results indicate a strong bottom-up control of functionally important herbivores mediated by woody plant phylogenetic diversity, and they show no sign of the increase in top-down effects of predator biomass or abundance generally expected to occur with increasing plant diversity (Root 1973; Haddad et al. 2009). Our findings have important implications for the regulation of ecosystem functions via trophic interaction pathways in these species-rich forests.

Plant species richness vs. phylogenetic diversity effects

While the comparatively weak direct influence of plant species richness on arthropod biomass and abundance in part contrasts with the findings from some other systems (mostly experimental grasslands; e.g. Haddad et al. 2009; Scherber et al. 2010), many other studies have likewise found no, or only weak and inconsistent, plant species richness effects (e.g. Koricheva et al. 2000; Perner et al. 2005; Vehviläinen et al. 2007; Vehviläinen et al. 2008). Interestingly, plant species richness effects are often considered to be mediated, and outperformed, by the influence of plant species composition (Perner et al. 2005; Zhang and Adams 2011; Rzanny et al. 2013). Metrics of plant diversity that account for the relatedness among species may thus help to reconcile contrasting findings by more effectively revealing the complexity of plant diversity effects beyond the mere impact of plant species richness (see Dinnage et al. 2012). This is demonstrated in our study, and for the first time for such species-rich forests, by the strong effects of woody plant phylogenetic diversity compared to overall weaker plant species richness effects. Many herbivores show phylogenetically structured host use (e.g. Weiblen et al. 2006), and phylodiversity-dependent patterns in herbivore assemblages might cascade to affect particularly specialized predators, whereas the response of generalist predators might be less pronounced. Thus, it is not surprising that the effect of woody plant phylogenetic diversity was particularly evident for herbivores and less so for predators, which in our case were largely generalist spiders. However, interactive effects of plant species richness and phylogenetic diversity on the predator to herbivore biomass ratio and on the biomass dispersion of leaf chewers indicate that species richness can contain information that is not fully captured by phylogenetic diversity, as can be the case when functionally important traits are not phylogenetically conserved (Srivastava et al. 2012; Schuldt et al. 2014b).

Leaf-chewing vs. sap-sucking herbivores

Leaf-chewing herbivores are responsible for the majority of visible leaf damage in the studied forest (Schuldt et al. 2010; Schuldt et al. 2012), and they also represented the largest proportion of the arthropod biomass in our samples (ca. 57% of the total and 85% of the herbivore biomass). The increase in leaf chewer biomass and abundance with higher woody plant phylogenetic diversity corresponds well to the increase in herbivore damage with increasing woody plant diversity found previously on the same study plots (Schuldt et al. 2014a). These patterns strongly suggest that the functional impact of herbivores increases with woody plant diversity. This contrasts with the general expectations of the resource-concentration hypothesis (Root 1973; Haddad et al. 2009) and also with commonly held assumptions that herbivores become increasingly specialized toward lower latitudes (Coley and Barone 1996; Dyer et al. 2007). However, our previous studies provided strong indications that the dominant herbivores in this system are generalists that may benefit from higher resource diversity in the more plant-diverse forest stands (Schuldt et al. 2014a). This mechanism, formulated by the ‘dietary-mixing’ hypothesis (Bernays et al. 1994; Dinnage 2013), and the probable dominance of generalist herbivores (which has also been suggested for other species-rich forests; e.g. Novotny et al. 2012), may explain the observed positive relationship between leaf chewer biomass and woody plant phylogenetic diversity. In particular, if the defense or palatability traits most relevant for herbivores show phylogenetic clustering, phylogenetically more diverse plant communities provide generalist herbivores with alternative hosts that help overcome dietary limitations on herbivore performance (Cavender-Bares et al. 2009; see Parker et al. 2012 for a real-world example). Leaf traits potentially important to herbivores that were found (among a set of 21 species) to show a phylogenetic signal at our study sites were e.g. dry matter content, toughness, polyphenols, tannins, and carbon and nitrogen contents (Schuldt et al. 2012; Eichenberg et al. 2014). Interactions among such traits and responses of individual herbivore species from the multidiverse set of herbivores potentially attacking a given plant species may make the

functional response of plants highly complex, and our phylogenetic diversity metric might capture the overall response of herbivore assemblages by integrating over the evolutionary adaptations of these herbivores. Without doubt, however, plant functional traits need to be explicitly addressed to identify the mechanisms underlying the observed diversity effects, and traits might also provide additional information on variation in the arthropod data not explained in our current models (see also Schuldt et al. 2014a). For instance, the fact that in some cases we observed effects of woody plant species richness beyond those of plant phylogenetic diversity might be a signal of functional trait information that is not phylogenetically conserved. The multitude of chemical, morphological and physiological traits that might potentially play a role in affecting arthropods, and the complexity of potential relationships among traits, however, require an extensive trait dataset for further testing that is often not available for many study regions. Yet, we hope that the results of our study will help to motivate efforts to unveil the complexity behind trophic interactions that might be key for ecosystem functioning in many species-rich systems.

The assumption that resource-diversity effects and not an increase in plant productivity were underlying reasons for the effects of plant phylogenetic diversity is supported by two additional findings of our study. First of all, even though primary productivity was previously found to increase with woody plant diversity (Baruffol et al. 2013), in our study it was not significantly related to any of the arthropod biomass and abundance patterns that showed a relationship with woody plant phylogenetic diversity. While the impact of plant diversity on arthropods operating through an increase in plant biomass may be common (e.g. Borer et al. 2012), the strength of these effects can vary and they may be overruled by resource diversity effects (Perner et al. 2005; Haddad et al. 2009; Dinnage et al. 2012). Secondly, the biomass distribution of leaf chewers within plots (Q_{Bio}) became more diverse with increasing plant phylogenetic diversity and woody plant species richness, which might be indicative of increased niche separation among herbivores due to a more

heterogeneous resource distribution (Mason et al. 2005). This higher biomass diversity, in turn, may contribute to strengthening the previously observed top-down effects of herbivores on the producer level with increasing plant diversity in the studied forest stands (see also Rudolf 2012).

The increase in herbivore biomass and abundance which we found differs from recent studies in grasslands that reported no significant effects of plant phylogenetic diversity on herbivore abundance patterns (in contrast to stronger effects on herbivore species richness; Dinnage et al. 2012; Pellissier et al. 2013). However, these studies indicated either a strong top-down control of herbivore abundances by predators (Dinnage et al. 2012) or focused on specialized herbivore taxa (Pellissier et al. 2013). As our results show, predator top-down control seems to be, at best, weak for the arthropod assemblages of woody plant saplings at our study site, and the probable dominance of generalist herbivores may explain deviations from patterns for more specialized taxa.

Such deviations between herbivore taxa also became evident to some extent in our study, as we found no significant effect of plant phylogenetic diversity on the biomass of sap-sucking herbivores. Indeed, we did not find sap sucker biomass to be related to any of the plot characteristics. And while the biomass dispersion of sap suckers increased with plant phylogenetic diversity in summer (Time 3), this relationship was not evident in spring and fall, again suggesting an overall much weaker effect of plant diversity on sap suckers as compared to leaf chewers. Differences in feeding mode and in the degree of host specialization could have caused these feeding guild-specific results, but the ultimate drivers are difficult to elucidate with our study. Nevertheless, several studies that included sap suckers in their analyses likewise found no or only weak direct effects of plant diversity measures on these herbivores (e.g. Koricheva et al. 2000; Unsicker et al. 2006; Vehviläinen et al. 2007). As sap suckers pierce plants to consume assimilates from phloem, xylem, or individual cells, they trigger different signaling pathways in plants and are able to avoid many

of the morphological and chemical defense mechanisms that deter leaf chewers (Howe and Jander 2008; Zvereva et al. 2010) and that may be related to the phylogenetic structure and diversity of plant communities (Baraloto et al. 2012). This may weaken potential relationships with plant phylogenetic diversity for sap-sucking herbivores and explain the patterns we found in our study, especially if a dominance of leaf chewers shifts plant responses to a stronger defense against these dominant herbivores (see also Carmona and Fornoni 2013).

Analyses at the level of individual arthropod species might potentially provide further insight, but are beyond the scope of our study. Moreover, as most herbivores show at least some degree of lineage-specificity in their host use (Weiblen et al. 2006), and considering that we sampled a wide range of woody plant species in each of our plots and analyzed mean values for arthropods averaged across all woody plant individuals sampled per plot, it is unlikely that our results are driven by the response of only a few specific arthropod species from specific woody plant species. Rather, the overall changes in biomass point to more general effects across larger parts of the herbivore assemblages. This is supported by our finding that the biomass distribution (Q_{Bio}) of leaf chewers increased with woody plant phylogenetic diversity, indicating that a wider range of species of different body size were promoted.

Predators and the predator-herbivore ratio

In contrast to herbivores feeding directly on plants, organisms at higher trophic levels are less strongly related to the plant community. The impact of plant diversity may thus be expected to become weaker higher up in food webs (Scherber et al. 2010). However, several recent studies have shown that predator abundance can strongly increase with plant diversity and particularly also with plant phylogenetic diversity (e.g. Haddad et al. 2009; Dinnage et al. 2012). This may occur either through positive bottom-up effects of the quantity of available prey resources or via increased structural and non-trophic components of more diverse plant

communities (Root 1973; Haddad et al. 2009). While our study indicates that predator abundance (but not biomass) was positively related to herbivore abundance, the strong effects of plant phylogenetic diversity on the latter did not translate to the predator level. The lack of plant phylogenetic diversity effects might be due to the fact that generalist predators (predominantly spiders), which may be less responsive to effects of plant phylogeny on herbivore assemblage structure than specialized predators, made up the largest proportion of both total predator biomass and abundance. Moreover, considering that the diet of generalist predators is made up of various herbivore guilds, the lack of a plant phylogenetic diversity effect on sap suckers may have contributed to suppressing an overall plant diversity effect on predators.

Overall, these patterns indicate that predators exert weak top-down control on herbivore biomass and abundance in the undergrowth of the studied forest stands, and that bottom-up effects of the producer level prevail. This is also supported by the decreasing ratios of predator to herbivore biomass and abundance with increasing plant phylogenetic diversity and woody plant species richness. With regard to plant diversity-dependent regulation mechanisms of ecosystem functions, this suggests that the often hypothesized strong mediating role of higher trophic levels for such species-rich forests (Schemske et al. 2009; Terborgh 2012) in our case particularly applies to dominant herbivores and less so to predators or less abundant herbivores. As our study focused on tree and shrub recruits rather than on the established canopy tree community, these patterns may have strong effects on the long-term structuring of the woody plant communities. Our previous studies showed that saplings of abundant tree and shrub species experienced greater damage than less common species and that this damage increased with woody plant diversity (Schuldt et al. 2010; Schuldt et al. 2012). The higher biomass and abundance of herbivores in the more diverse forest stands may thus actually contribute to maintaining this high plant diversity, by decreasing the performance of common tree and shrub species, and thus promoting

coexistence with less common species (see also Dyer et al. 2010).

Conclusions

The strong effects of woody plant phylogenetic diversity, and the much less pronounced direct effects of plant species richness, on herbivore biomass and abundance show that the diversity-dependence of ecosystem processes and interaction pathways across trophic levels can fundamentally depend on non-random associations among species. Scenarios of random species loss may thus underestimate the consequences for ecosystem functions if they do not reflect the driving forces of community assembly (see also Dinnage et al. 2012). Our study particularly highlights the impact of strong bottom-up effects of plant phylogenetic diversity, whereas top-down effects of predators, at least for the undergrowth of tree and shrub recruits studied here, are less responsive to plant diversity and in this context exert weak effects on herbivores. However, this means that herbivores may, in turn, strongly impact particular plant species, which may potentially lead to a positive feedback loop of bottom-up controlled herbivores on plant diversity maintenance. Moreover, considering that biomass is directly related to metabolic rates and the performance of individuals (Saint-Germain et al. 2007; Reiss et al. 2011), the more pronounced effects of plant diversity on arthropod biomass than on abundance found in our study suggest that abundance patterns alone (although more frequently analyzed than biomass patterns in ecological studies; Reiss et al. 2011) may be less effective in capturing the diversity-dependent impact of arthropods in such species-rich ecosystems.

Acknowledgements

We thank the administration of the Gutianshan National Nature Reserve and members of the BEF China consortium for support, Stefan Michalski and Walter Durka for providing woody plant phylogenetic data, and Daria Golub, Merle Noack, and Friederike Rorig for help with

arthropod sampling. Comments of two anonymous reviewers greatly helped to improve the manuscript. We gratefully acknowledge funding by the German Research Foundation (DFG FOR 891/1 and 891/2), the Sino-German Centre for Research Promotion in Beijing (GZ 524, 592, 698, 699, 785, and 1020), and the National Science Foundation of China (NSFC 30710103907 and 309300005).

References

- Baraloto C et al. (2012) Using functional traits and phylogenetic trees to examine the assembly of tropical tree communities. *J Ecol* 100:690-701
- Baruffol M et al. (2013) Biodiversity promotes tree growth during succession in subtropical forest. *PLoS ONE* 8:e81246
- Bernays EA, Bright KL, Gonzalez N, Angel J (1994) Dietary mixing in a generalist herbivore: tests of two hypotheses. *Ecology* 75:1997-2006
- Borer ET, Seabloom EW, Tilman D (2012) Plant diversity controls arthropod biomass and temporal stability. *Ecol Lett* 15:1457-1464
- Botta-Dukát Z (2005) Rao's quadratic entropy as a measure of functional diversity based on multiple traits. *J Veg Sci* 16:533-540
- Bruehlheide H et al. (2011) Community assembly during secondary forest succession in a Chinese subtropical forest. *Ecol Monogr* 81:25-41
- Burnham KP, Anderson DR (2004) Model selection and multimodel inference : a practical information-theoretic approach. Springer, New York
- Campos RI, Vasconcelos HL, Ribeiro SP, Neves FS, Soares JP (2006) Relationship between tree size and insect assemblages associated with *Anadenanthera macrocarpa*. *Ecography* 29:442-450
- Cardinale BJ et al. (2012) Biodiversity loss and its impact on humanity. *Nature* 486:59-67
- Cardinale BJ et al. (2011) The functional role of producer diversity in ecosystems. *Am J Bot*

540 98:572-592

541 Carmona D, Fornoni J (2013) Herbivores can select for mixed defensive strategies in plants.

542 New Phytol 197:576-585

543 Castagneyrol B, Jactel H (2012) Unraveling plant–animal diversity relationships: a meta-

544 regression analysis. Ecology 93:2115-2124

545 Cavender-Bares J, Kozak KH, Fine PVA, Kembel SW (2009) The merging of community

546 ecology and phylogenetic biology. Ecol Lett 12:693-715

547 Coley PD, Barone JA (1996) Herbivory and plant defenses in tropical forests. Annu Rev Ecol

548 Syst 27:305-335

549 Dinnage R (2013) Phylogenetic diversity of plants alters the effect of species richness on

550 invertebrate herbivory. PeerJ 1:e93

551 Dinnage R, Cadotte MW, Haddad NM, Crutsinger GM, Tilman D (2012) Diversity of plant

552 evolutionary lineages promotes arthropod diversity. Ecol Lett 15:1308-1317

553 Dyer LA, Letourneau DK, Chavarria GV, Amoretti DS (2010) Herbivores on a dominant

554 understory shrub increase local plant diversity in rain forest communities. Ecology

555 91:3707-3718

556 Dyer LA et al. (2007) Host specificity of Lepidoptera in tropical and temperate forests. Nature

557 448:696-U699

558 Eichenberg D, Ristok C, Kröber W, Bruelheide H (2014) Plant polyphenols - implications of

559 different sampling, storage and sample processing in biodiversity-ecosystem functioning

560 experiments. Chem Ecol. DOI: 10.1080/02757540.2014.894987

561 Futuyma DJ, Agrawal AA (2009) Macroevolution and the biological diversity of plants and

562 herbivores. Proc Natl Acad Sci USA 106:18054-18061

563 Geißler C, Kühn P, Böhnke M, Bruelheide H, Shi X, Scholten T (2012) Splash erosion

564 potential under tree canopies in subtropical SE China. Catena 91:85-93

565 Gotelli NJ, Rohde K (2002) Co-occurrence of ectoparasites of marine fishes: a null model

analysis. *Ecol Lett* 5:86-94

Haddad NM, Crutsinger GM, Gross K, Haarstad J, Knops JMH, Tilman D (2009) Plant species loss decreases arthropod diversity and shifts trophic structure. *Ecol Lett* 12:1029-1039

Hardy OJ (2008) Testing the spatial phylogenetic structure of local communities: statistical performances of different null models and test statistics on a locally neutral community. *J Ecol* 96:914-926

Hódar JA (1996) The use of regression equations for estimation of arthropod biomass in ecological studies. *Acta Oecol* 17:421-433

Hooper DU et al. (2012) A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature* 486:105-108

Howe GA, Jander G (2008) Plant immunity to insect herbivores. *Annu Rev Plant Biol* 59:41-66

Hu Z, Yu M (2008) Study on successions sequence of evergreen broad-leaved forest in Gutian Mountain of Zhejiang, Eastern China: species diversity. *Front Biol China* 3:45-49

Kembel SW et al. (2010) Picante: R tools for integrating phylogenies and ecology. *Bioinformatics* 26:1463-1464

Koricheva J, Mulder CPH, Schmid B, Joshi J, Huss-Danell K (2000) Numerical responses of different trophic groups of invertebrates to manipulations of plant diversity in grasslands. *Oecologia* 125:271-282

Legendre P et al. (2009) Partitioning beta diversity in a subtropical broad-leaved forest of China. *Ecology* 90:663-674

Mason NWH, Mouillot D, Lee WG, Wilson JB (2005) Functional richness, functional evenness and functional divergence: the primary components of functional diversity. *Oikos* 111:112-118

Nadrowski K, Wirth C, Scherer-Lorenzen M (2010) Is forest diversity driving ecosystem

592 functioning and service? *Curr Opin Envir Sus* 2:75-79

593 Naeem S, Duffy JE, Zavaleta E (2012) The functions of biological diversity in an age of
 594 extinction. *Science* 336:1401-1406

595 Novotny V et al. (2010) Guild-specific patterns of species richness and host specialization in
 596 plant-herbivore food webs from a tropical forest. *J Anim Ecol* 79:1193-1203

597 Novotny V et al. (2012) Insects on plants: explaining the paradox of low diversity within
 598 specialist herbivore guilds. *Am Nat* 179:351-362

599 Ødegaard F, Diserud OH, Østbye K (2005) The importance of plant relatedness for host
 600 utilization among phytophagous insects. *Ecol Lett* 8:612-617

601 Paine CET et al. (2012) Phylogenetic density dependence and environmental filtering predict
 602 seedling mortality in a tropical forest. *Ecol Lett* 15:34-41

603 Parker JD, Burkepille DE, Lajeunesse MJ, Lind EM (2012) Phylogenetic isolation increases
 604 plant success despite increasing susceptibility to generalist herbivores. *Divers Distrib*
 605 18:1-9

606 Pellissier L et al. (2013) Turnover of plant lineages shapes herbivore phylogenetic beta
 607 diversity along ecological gradients. *Ecol Lett* 16:600-608

608 Perner J et al. (2005) Effects of plant diversity, plant productivity and habitat parameters on
 609 arthropod abundance in montane European grasslands. *Ecography* 28:429-442

610 Pinheiro J, Bates D, Debroy S, Sarkar D, R Core Team (2014) nlme: linear and nonlinear
 611 mixed effects models. R package version 3.1-117, available at [http://CRAN.R-](http://CRAN.R-project.org/package=nlme)
 612 [project.org/package=nlme](http://CRAN.R-project.org/package=nlme). Last accessed 26 May 2014.

613 Plath M, Dorn S, Riedel J, Barrios H, Mody K (2012) Associational resistance and
 614 associational susceptibility: specialist herbivores show contrasting responses to tree stand
 615 diversification. *Oecologia* 169:477-487

616 Purschke O, Michalski SG, Bruelheide H, Durka W (2014) Spatial and temporal phylogenetic
 617 turnover during subtropical forest succession across ecological and phylogenetic scales. *J*

618 Ecol:in review

619 Reiss J, Bailey RA, Perkins DM, Pluchinotta A, Woodward G (2011) Testing effects of
620 consumer richness, evenness and body size on ecosystem functioning. *J Anim Ecol*
621 80:1145-1154

622 Root RB (1973) Organization of a plant-arthropod association in simple and diverse habitats:
623 the fauna of collards (*Brassica oleracea*). *Ecol Monogr* 43:95-124

624 Rudolf VHW (2012) Seasonal shifts in predator body size diversity and trophic interactions in
625 size-structured predator–prey systems. *J Anim Ecol* 81:524-532

626 Rzanny M, Kuu A, Voigt W (2013) Bottom-up and top-down forces structuring consumer
627 communities in an experimental grassland. *Oikos* 122:967-976

628 Saint-Germain M et al. (2007) Should biomass be considered more frequently as a currency in
629 terrestrial arthropod community analyses? *J Appl Ecol* 44:330-339

630 Schemske DW, Mittelbach GG, Cornell HV, Sobel JM, Roy K (2009) Is there a latitudinal
631 gradient in the importance of biotic interactions? *Annu Rev Ecol Evol Syst* 40:245-269

632 Scherber C et al. (2010) Bottom-up effects of plant diversity on multitrophic interactions in a
633 biodiversity experiment. *Nature* 468:553-556

634 Schleuter D, Daufresne M, Massol F, Argillier C (2010) A user's guide to functional diversity
635 indices. *Ecol Monogr* 80:469-484

636 Schuldt A et al. (2014a) Functional and phylogenetic diversity of woody plants drive
637 herbivory in a highly diverse forest. *New Phytol* 202:864-873

638 Schuldt A et al. (2010) Tree diversity promotes insect herbivory in subtropical forests of
639 south-east China. *J Ecol* 98:917-926

640 Schuldt A et al. (2012) Plant traits affecting herbivory on tree recruits in highly diverse
641 subtropical forests. *Ecol Lett* 15:732-739

642 Schuldt A, Bruelheide H, Durka W, Michalski SG, Purschke O, Assmann T (2014b) Tree
643 diversity promotes functional dissimilarity and maintains functional richness despite

644 species loss in predator assemblages. *Oecologia* 174:533-543

645 Sobek S, Scherber C, Steffan-Dewenter I, Tschardt T (2009a) Sapling herbivory,
646 invertebrate herbivores and predators across a natural tree diversity gradient in Germany's
647 largest connected deciduous forest. *Oecologia* 160:279-288

648 Sobek S, Steffan-Dewenter I, Scherber C, Tschardt T (2009b) Spatiotemporal changes of
649 beetle communities across a tree diversity gradient. *Divers Distrib* 15:660-670

650 Srivastava DS, Cadotte MW, MacDonald AAM, Marushia RG, Mirotchnick N (2012)
651 Phylogenetic diversity and the functioning of ecosystems. *Ecol Lett* 15:637-648

652 Srivastava DS, Lawton JH (1998) Why more productive sites have more species: An
653 experimental test of theory using tree-hole communities. *Am Nat* 152:510-529

654 Srivastava DS, Vellend M (2005) Biodiversity-ecosystem function research: Is it relevant to
655 conservation? *Annu Rev Ecol Evol Syst* 36:267-294

656 Strutzenberger P, Brehm G, Fiedler K (2011) DNA barcoding-based species delimitation
657 increases species count of Eois (Geometridae) moths in a well-studied tropical mountain
658 forest by up to 50%. *Insect Sci* 18:349-362

659 Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S (2011) MEGA5: Molecular
660 evolutionary genetics analysis using maximum likelihood, evolutionary distance, and
661 maximum parsimony methods. *Mol Biol Evol* 28:2731-2739

662 Terborgh J (2012) Enemies maintain hyperdiverse tropical forests. *Am Nat* 179:303-314

663 Thebault E, Huber V, Loreau M (2007) Cascading extinctions and ecosystem functioning:
664 contrasting effects of diversity depending on food web structure. *Oikos* 116:163-173

665 Unsicker SB, Baer N, Kahmen A, Wagner M, Buchmann N, Weisser WW (2006) Invertebrate
666 herbivory along a gradient of plant species diversity in extensively managed grasslands.
667 *Oecologia* 150:233-246

668 Vehviläinen H, Koricheva J, Ruohomäki K (2007) Tree species diversity influences herbivore
669 abundance and damage: meta-analysis of long-term forest experiments. *Oecologia*

670 152:287-298

671 Vehviläinen H, Koricheva J, Ruohomäki K (2008) Effects of stand tree species composition
672 and diversity on abundance of predatory arthropods. *Oikos* 117:935-943

673 Wardhaugh CW (2013) Estimation of biomass from body length and width for tropical
674 rainforest canopy invertebrates. *Aust J Entomol* 52:291-298

675 Wardhaugh CW, Stork NE, Edwards W (2012) Feeding guild structure of beetles on
676 Australian tropical rainforest trees reflects microhabitat resource availability. *J Anim*
677 *Ecol* 81:1086-1094

678 Weiblen GD, Webb CO, Novotny V, Basset Y, Miller SE (2006) Phylogenetic dispersion of
679 host use in a tropical insect herbivore community. *Ecology* 87:S62-S75

680 Whitfeld TJS, Novotny V, Miller SE, Hrcek J, Klimes P, Weiblen GD (2012) Predicting
681 tropical insect herbivore abundance from host plant traits and phylogeny. *Ecology*
682 93:S211-S222

683 Zhang Y, Adams J (2011) Top-down control of herbivores varies with ecosystem types. *J*
684 *Ecol* 99:370-372

685 Zvereva E, Lanta V, Kozlov M (2010) Effects of sap-feeding insect herbivores on growth and
686 reproduction of woody plants: a meta-analysis of experimental studies. *Oecologia*
687 163:949-960

Table 1. Mixed-effects models for the biomass of leaf-chewing herbivores, sap-sucking herbivores, predators, and the biomass ratio of predators to herbivores across 27 forest stands in subtropical China. Parameter estimates (with standard errors, degrees of freedom, t and *P* values) are shown for the variables retained in the minimal models

<i>Fixed effects</i> ^{a,b}	Herbivores: Leaf chewers					Herbivores: Suckers					Predators					Predator : Herbivore ratio				
	Est.	SE	DF	t	P	Est.	SE	DF	t	P	Est.	SE	DF	t	P	Est.	SE	DF	t	P
Intercept	1.53	0.14	52	10.6	<0.001	-0.64	0.23	52	-2.8	0.007	1.40	0.06	54	23.0	<0.001	-0.40	0.16	52	-2.5	0.016
Time 2-1 ^c	0.61	0.19	52	3.2	0.003	1.40	0.32	52	4.3	<0.001	-	-	-	-	-	-0.47	0.22	52	-2.2	0.035
Time 3-1 ^c	0.26	0.19	52	1.3	0.183	-0.51	0.32	52	-1.6	0.122	-	-	-	-	-	-0.05	0.22	52	-0.2	0.825
Plot age	-	-	-	-	-	-	-	-	-	-	0.17	0.06	25	2.8	0.009	-	-	-	-	-
Woody plant species richness (log)	0.18	0.10	24	1.9	0.067	-	-	-	-	-	-	-	-	-	-	-0.18	0.10	23	-1.8	0.093
<i>Qphyl</i> s.e.s. ^c	0.21	0.10	24	2.3	0.033	-	-	-	-	-	-	-	-	-	-	-0.28	0.12	23	-2.4	0.026
Woody plant richness : <i>Qphyl</i> s.e.s.	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.30	0.13	23	2.4	0.025
AICc full model ^d	205.4					298.1					153.9					231.4				
AICc min. model ^d	192.4					266.0					127.5					206.9				

^aSapling height, elevation, and the interactions *Qphyl* s.e.s : plot age, richness : plot age, time : *Qphyl* s.e.s , time : plot age, and time : richness were included in the full models but not retained in any of the minimal models

^bItalics denote data for non-significant terms retained in the minimal models

^c*Qphyl* s.e.s. = standardized effect sizes of woody plant phylogenetic diversity (*Qphyl*)

^dAkaike information criterion (corrected for small sample sizes) of the full model (containing all predictors) and the minimal adequate model (simplified model with lowest AICc)

Table 2. Mixed-effects models for the functional dispersion (Q_{Bio}) of the biomass of leaf chewer, sap-sucker, and predator individuals per plot across 27 forest stands in subtropical China. Parameter estimates (with standard errors, degrees of freedom, t and P values) are shown for the variables retained in the minimal models

	Herbivores: Leaf chewers					Herbivores: Suckers					Predators				
<i>Fixed effects</i> ^{a,b}	Est.	SE	DF	t	P	Est.	SE	DF	t	P	Est.	SE	DF	t	P
Intercept	-3.58	0.05	54	-75.4	<0.001	-4.25	0.21	50	-20.5	<0.001	-2.28	0.03	50	-72.2	<0.001
Time 2-1 ^c	-	-	-	-	-	0.73	0.29	50	2.5	0.016	0.07	0.04	50	1.5	0.148
Time 3-1 ^c	-	-	-	-	-	0.43	0.29	50	1.5	0.149	0.18	0.04	50	4.0	<0.001
Plot age	0.09	0.05	22	2.0	0.058	-	-	-	-	-	-	-	-	-	-
Woody plant species richness															
(log)	0.12	0.05	22	2.4	0.024	-	-	-	-	-	0.05	0.03	25	1.4	0.166
<i>Qphyl</i> s.e.s.	0.19	0.06	22	3.4	0.003	-0.06	0.21	25	-0.3	0.759	-	-	-	-	-
Woody plant richness : <i>Qphyl</i>															
s.e.s.	-0.15	0.06	22	-2.5	0.022	-	-	-	-	-	-	-	-	-	-
Time 2 : <i>Qphyl</i>															
s.e.s.	-	-	-	-	-	-0.04	0.29	50	-0.1	0.890	-	-	-	-	-
Time 3 : <i>Qphyl</i>															
s.e.s.	-	-	-	-	-	0.67	0.29	50	2.3	0.027	-	-	-	-	-

Time 2 : Woody																
plant richness	-	-	-	-	-	-	-	-	-	-	-0.11	0.04	50	-2.5	0.014	
Time 3 : Woody																
plant richness	-	-	-	-	-	-	-	-	-	-	-0.07	0.04	50	-1.6	0.108	
AICc full model	116.5						281.5				-24.4					
AICc min. model	88.4						253.1				-51.2					

^aSapling height, elevation, and the interactions *Qphyl* s.e.s : plot age, richness : plot age, and time : plot age were included in the full models but not retained in any of the minimal models

^bItalics denote data for non-significant terms retained in the minimal models

^c*Qphyl* s.e.s. = standardized effect sizes of woody plant phylogenetic diversity (*Qphyl*)

^dContrasts between sampling time 1 (fall) and the successive sampling times 2 and 3 (spring and summer); the intercept is the overall mean

^eAkaike information criterion (corrected for small sample sizes) of the full model (containing all predictors) and the minimal adequate model (simplified model with lowest AICc)




Figure 1. Relationships between woody plant phylogenetic diversity (Q_{phyl} s.e.s.), woody plant species richness, plot age and the biomass of a) leaf chewers, b) predators, and c) the ratio of predators to herbivores across 27 forest plots in subtropical China. All relationships are significant at $P < 0.05$ (see Table 1 for details).




Figure 2. Relationships between the dispersion of biomass per plot among individuals of leaf-chewing herbivores and woody plant phylogenetic diversity (Q_{phyl} s.e.s) and woody plant species richness. All relationships are significant at $P < 0.05$ (Table 1).